



PROGRAMME T SALMO

Vers. 30/10/2015

Table de nourrissage indicative (kg d'aliment / 100 kg de biomasse / jour)

	PROT. (%)	LIP. (%)	ED (MJ/kg)	PD/ED (g/MJ)	Diamètre		Poids vifs (g)		Température de l'eau																
					n°	(mm)	de	à	4°C	5°C	6°C	7°C	8°C	9°C	10°C	11°C	12°C	13°C	14°C	15°C	16°C	17°C	18°C	19°C	20°C
T SALMO 19	45	21	19.4	21.4	3	3.2	10	50	1.03	1.1	1.2	1.27	1.35	1.44	1.52	1.6	1.7	1.77	1.82	1.85	1.86	1.84	1.8	1.68	1.52
	42	24	19.7	19.2	4	4	50	100	0.9	0.97	1.05	1.13	1.2	1.3	1.38	1.46	1.53	1.6	1.65	1.7	1.72	1.7	1.64	1.5	1.38
	40	25	19.7	18.2	5	5.5	100	150	0.82	0.9	0.97	1.05	1.12	1.2	1.27	1.36	1.43	1.5	1.57	1.6	1.62	1.6	1.52	1.4	1.27
					5	5.5	150	200	0.8	0.85	0.92	1	1.07	1.13	1.2	1.27	1.35	1.42	1.5	1.52	1.52	1.5	1.42	1.32	1.2
					5	5.5	200	400	0.72	0.77	0.84	0.9	0.98	1.05	1.12	1.17	1.24	1.3	1.37	1.4	1.4	1.37	1.3	1.22	1.12
T SALMO 20	39	27	20.4	17.2	7	7.5	400	500	0.64	0.7	0.75	0.82	0.87	0.94	1	1.06	1.12	1.18	1.23	1.25	1.25	1.23	1.18	1.1	1
					7	7.5	500	1000	0.58	0.63	0.68	0.74	0.8	0.85	0.9	0.96	1	1.05	1.1	1.13	1.12	1.07	1	0.9	
					7	7.5	1000	1500	0.5	0.55	0.6	0.64	0.68	0.73	0.8	0.83	0.88	0.9	0.95	0.97	0.98	0.97	0.93	0.86	0.8
					7	7.5	1500	1800	0.46	0.5	0.54	0.58	0.63	0.67	0.7	0.75	0.8	0.83	0.87	0.9	0.9	0.88	0.84	0.78	0.7
T SALMO 21	37	30	20.9	16	7	7.5	400	500	0.63	0.67	0.74	0.8	0.85	0.9	0.98	1.03	1.1	1.15	1.2	1.22	1.22	1.2	1.15	1.06	0.98
					7	7.5	500	1000	0.57	0.62	0.67	0.72	0.78	0.83	0.9	0.94	1	1.03	1.06	1.08	1.1	1.1	1.04	0.97	0.9
					7	7.5	1000	1500	0.5	0.54	0.58	0.62	0.67	0.72	0.77	0.8	0.86	0.9	0.93	0.94	0.96	0.94	0.9	0.84	0.77
					7	7.5	1500	1800	0.45	0.5	0.53	0.57	0.6	0.65	0.7	0.73	0.78	0.8	0.85	0.87	0.87	0.86	0.82	0.77	0.7
					9	9.5	1800	2000	0.43	0.47	0.5	0.54	0.58	0.62	0.66	0.7	0.73	0.77	0.8	0.82	0.82	0.8	0.78	0.72	0.66
					9	9.5	2000	2500	0.4	0.43	0.46	0.5	0.53	0.57	0.6	0.64	0.67	0.7	0.74	0.77	0.76	0.77	0.76	0.72	0.67
T SALMO 22	36	32	21.7	15.3	9	9.5	1800	2000	0.4	0.45	0.5	0.52	0.56	0.6	0.63	0.67	0.7	0.74	0.77	0.8	0.8	0.78	0.75	0.7	0.63
					9	9.5	2000	2500	0.38	0.4	0.45	0.48	0.5	0.55	0.58	0.62	0.65	0.7	0.72	0.73	0.74	0.73	0.7	0.65	0.58
					11	11	2500	3000	0.35	0.38	0.4	0.43	0.47	0.5	0.52	0.55	0.6	0.6	0.64	0.65	0.65	0.65	0.6	0.57	0.52
					11	11	3000	3500	0.3	0.33	0.36	0.4	0.4	0.44	0.47	0.5	0.52	0.55	0.57	0.6	0.6	0.58	0.55	0.5	0.47
					11	11	3500	4000	0.28	0.3	0.32	0.35	0.37	0.4	0.42	0.44	0.47	0.5	0.52	0.53	0.53	0.52	0.48	0.45	0.42

Les taux de nourrissage sont basés sur la valeur nutritionnelle de l'aliment et doivent être adaptés aux conditions locales des élevages.

